

From J. M. Thijssen, Computational Physics (Cambridge University Press, 1999)

## 2

# Quantum scattering with a spherically symmetric potential

## 2.1 Introduction

In this chapter, we shall discuss quantum scattering with a spherically symmetric potential as a typical example of the problems studied in computational physics.<sup>1,6</sup> Scattering experiments are perhaps the most important tool for obtaining detailed information on the structure of matter, in particular the interaction between particles. Examples of scattering techniques include neutron and X-ray scattering for liquids, atoms scattering from crystal surfaces, elementary particle collisions in accelerators. In most of these scattering experiments, a beam of incident particles hits a target which also consists of many particles. The distribution of scattering particles over the different directions is then measured, for different energies of the incident particles. This distribution is the result of many individual scattering events. Quantum mechanics enables us, in principle, to evaluate for an individual event the probabilities for the incident particles to be scattered off in different directions; and this probability is identified with the measured distribution.

Suppose we have an idea of what the potential between the particles involved in the scattering process might look like, for example from quantum mechanical energy calculations (programs for this purpose will be discussed in the next few chapters). We can then *parametrise* the interaction potential, i.e. we write it as an analytic expression involving a set of constants: the parameters. If we evaluate the scattering probability as a function of the scattering angles for different values of these parameters, and compare the results with experimental scattering data, we can find those parameter values for which the agreement between theory and experiment is optimal. Of course, it would be nice if we could evaluate the scattering potential directly from the scattering data (this is called the *inverse problem*), but this is unfortunately




Figure 2.1: Geometry of a scattering process.

very difficult (if not impossible) as many different interaction potentials can have similar scattering properties as we shall see below.

Many different motivations for obtaining accurate interaction potentials can be given. One is that we might use the interaction potential to make predictions about the behaviour of a system consisting of many interacting particles, such as a dense gas or a liquid. Methods for doing this will be discussed in chapters 8 and 10.

Scattering might be *elastic* or *inelastic*. In the former case the energy is conserved, in the latter energy disappears. This means that energy transfer takes place from the scattered particles to degrees of freedom which are not included explicitly in the system (inclusion of these degrees of freedom would cause the energy to be conserved). In this chapter we shall consider elastic scattering. We restrict ourselves furthermore to spherically symmetric interaction potentials. In chapter 13 we shall briefly discuss scattering in the context of quantum field theory for elementary particles.

We analyse the scattering process of a particle incident on a scattering centre which is usually another particle.<sup>†</sup> We assume that we know the scattering potential which is spherically symmetric so that it depends on the distance between the particle and the scattering centre only.

In an experiment, one typically measures the scattered flux, that is the intensity of the outgoing beam for various directions which are denoted by the spatial angle  $\Omega = (\theta, \phi)$  as in figure 2.1. The *differential cross section*,  $\frac{d\sigma}{d\Omega}(\Omega)$ , describes how these intensities are distributed over the various spatial angles  $\Omega$  and the integrated flux of the scattered particles is the *total cross section*,  $\sigma_{\text{tot}}$ . These experimental quantities are what we want to calculate.

<sup>†</sup>Every two-particle collision can be transformed into a single scattering problem involving the relative position; in the transformed problem the incoming particle has the reduced mass  $m = m_1 m_2 / (m_1 + m_2)$ .

The scattering process is described by the solutions of the single-particle Schrödinger equation involving the (reduced) mass  $m$ , the relative coordinate  $\mathbf{r}$  and the interaction potential  $V$  between the particle and the interaction centre:

$$\left[ -\frac{\hbar^2}{2m} \nabla^2 + V(r) \right] \psi(\mathbf{r}) = E\psi(\mathbf{r}). \quad (2.1)$$

This is a partial differential equation in three dimensions, which could be solved using the 'brute force' discretisation methods presented in appendix A, but exploiting the spherical symmetry of the potential, we can solve the problem in another, more elegant, way which, moreover, works much faster on a computer. More specifically, in section 2.3 we shall establish a relation between the *phase shift* and the scattering cross sections. In this section, we shall restrict ourselves to a description of the concept of phase shift and describe how it can be obtained from the solutions of the radial Schrödinger equation. The expressions for the scattering cross sections will then be used to build the computer program which is described in section 2.2.

For the potential  $V(r)$  we make the assumption that it vanishes for  $r$  larger than a certain value  $r_{\max}$ . In case we are dealing with an asymptotically decaying potential, we neglect contributions from the potential beyond the range  $r_{\max}$ , which must be chosen suitably, or treat the tail in a perturbative manner as described in problem 2.2.

For a spherically symmetric potential, the solution of the Schrödinger equation can always be written as

$$\psi(\mathbf{r}) = \sum_{l=0}^{\infty} \sum_{m=-l}^l A_{lm} \frac{u_l(r)}{r} Y_l^m(\theta, \phi) \quad (2.2)$$

where  $u_l$  satisfies the radial Schrödinger equation:

$$\left\{ \frac{\hbar^2}{2m} \frac{d^2}{dr^2} + \left[ E - V(r) - \frac{\hbar^2 l(l+1)}{2mr^2} \right] \right\} u_l(r) = 0. \quad (2.3)$$

Figure 2.2 shows the solution of the radial Schrödinger equation with  $l=0$  for a square well potential for various well depths – our discussion applies also to nonzero values of  $l$ . Outside the well, the solution  $u_l$  can be written as a linear combination of the two independent solutions  $j_l$  and  $n_l$ , the regular and irregular spherical Bessel functions. We write this linear combination in the particular form

$$u_l(r > r_{\max}) \propto kr [\cos \delta_l j_l(kr) - \sin \delta_l n_l(kr)]. \quad (2.4)$$

Figure

 $\delta_l$  is d  
for w  
spheri

which

We s  
wave  
sine  
T  
abou  
enab  
secti

## 2.1. Introduction

17




Figure 2.2: The radial wave functions for  $l=0$  for various square well potential depths.

$\delta_l$  is determined via a matching procedure at the well boundary. The motivation for writing  $u_l$  in this form follows from the asymptotic expansion for the spherical Bessel functions:

$$kr j_l(kr) \approx \sin(kr - l\pi/2) \quad (2.5a)$$

$$kr n_l(kr) \approx -\cos(kr - l\pi/2) \quad (2.5b)$$

$$k = \sqrt{2mE}/\hbar$$

which can be used to rewrite (2.4) as

$$u_l(r) \propto \sin(kr - l\pi/2 + \delta_l), \quad \text{large } r. \quad (2.6)$$

We see that  $u_l$  approaches a sine-wave form for large  $r$  and the phase of this wave is determined by  $\delta_l$ , hence the name 'phase shift' for  $\delta_l$  (for  $l=0$   $u_l$  is a sine wave for all  $r > r_{\max}$ ).

The phase shift as a function of energy and  $l$  contains all the information about the scattering properties of the potential. In particular, the phase shift enables us to calculate the scattering cross sections and this will be done in section 2.3; here we simply quote the results. The differential cross section is

$$(2.4)$$

given in terms of the phase shift by

$$\frac{d\sigma}{d\Omega} = \frac{1}{k^2} \left| \sum_{l=0}^{\infty} (2l+1) e^{i\delta_l} \sin(\delta_l) P_l(\cos\theta) \right|^2 \quad (2.7)$$

and for the total cross section we find

$$\sigma_{\text{tot}} = 2\pi \int d\theta \sin\theta \frac{d\sigma}{d\Omega}(\theta) = \frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) \sin^2 \delta_l. \quad (2.8)$$

Summarising the analysis up to this point, we see that the potential determines the phase shift through the solution of the Schrödinger equation for  $r < r_{\text{max}}$ . The phase shift acts as an intermediate object between the interaction potential and the experimental scattering cross sections, as the latter can be determined from it.

Unfortunately, the expressions (2.7) and (2.8) contain sums over an infinite number of terms – hence they cannot be evaluated on the computer exactly. However, cutting off these sums can be motivated by a physical argument. Classically, only the waves with an angular momentum smaller than  $\hbar l_{\text{max}} = \hbar k r_{\text{max}}$  will ‘feel’ the potential – particles with higher  $l$ -values will pass by unaffected. Therefore we can safely cut off the sums at a somewhat higher value of  $l$  – we can always check whether the results obtained change significantly when taking more terms into account. We shall frequently encounter procedures similar to the cutting off described here. It is the art of computational physics to cleverly reduce infinite problems to ones which fit into the computer and still provide a reliable description.

How is the phase shift determined in practice? First, the Schrödinger equation must be integrated from  $r = 0$  outwards with boundary condition  $u_l(r=0) = 0$ . At  $r_{\text{max}}$ , the numerical solution must be matched onto the form (2.4) to fix  $\delta_l$ . The matching can be done either via the logarithmic derivative or using the value of the numerical solution at two different points  $r_1$  and  $r_2$  beyond  $r_{\text{max}}$  and we will use the latter method in order to avoid calculating derivatives. From (2.4) it follows directly that the phase shift is given by

$$\tan \delta_l = \frac{K j_l^{(1)} - j_l^{(2)}}{K n_l^{(1)} - n_l^{(2)}} \quad \text{with} \quad (2.9a)$$

$$K = \frac{r_1 u_l^{(2)}}{r_2 u_l^{(1)}}. \quad (2.9b)$$

In this equation,  $j_l^{(1)}$  stands for  $j_l(kr_1)$  etc.

## 2.2 A program for calculating cross sections

(2.7)

In this section we describe the construction of a program for calculating cross sections for a particular scattering problem: hydrogen atoms scattered off (much heavier) krypton atoms. Both atoms are considered as single particles and their structure (nucleus and electrons) is not explicitly taken into account. After completion, we are able to compare the results with experimental data. The program described here closely follows the work of Toennies *et al.* who carried out various atomic collisions experimentally and modelled the results using a similar computer program.<sup>7</sup>

The program is built up in several steps.

- First, the integration method for solving the radial Schrödinger equation is programmed. Various numerical methods can be used – we consider in particular Numerov's method (see section A.7.1.4).
- Second, we need routines yielding spherical Bessel functions in order to determine the phase shift via the matching procedure Eq. (2.9a). If we want to calculate differential cross sections, we need Legendre polynomials too. In section A.2, iterative methods for evaluating special functions are discussed.
- Finally, we complete the program with a routine for calculating the cross sections from the phase shifts.

### 2.2.1 Numerov's algorithm for the radial Schrödinger equation

The radial Schrödinger equation is given in Eq. (2.3). We define

$$F(l, r, E) = V(r) + \frac{\hbar^2 l(l+1)}{2mr^2} - E \quad (2.10)$$

so that the radial Schrödinger equation reads now:

$$\frac{\hbar^2}{2m} \frac{d^2}{dr^2} u(r) = F(l, r, E) u(r). \quad (2.11)$$

Units are chosen in which  $\hbar^2/(2m)$  assumes a reasonable value, that is, not extremely large and not extremely small – see below. You can choose a library routine for integrating this equation but if you prefer to write one yourself, Numerov's method is a good choice because it combines the simplicity of a regular mesh with good efficiency. The Runge-Kutta method can be used if

(2.9a)

(2.9b)

you want to have the freedom of varying the integration step when the potential changes rapidly – see problem 2.1.

Numerov's algorithm is described in section A.7.1.4. It makes use of the special structure of this equation to solve it with an error of order  $h^6$  ( $h$  is the discretisation interval) using only a three-point method. For  $\hbar^2/2m \equiv 1$  it reads:

$$w(r+h) = 2w(r) - w(r-h) + h^2 F(l, r, E) u(r) \quad (2.12)$$

and

$$u(r) = \left[ 1 - \frac{h^2}{12} F(l, r, E) \right]^{-1} w(r). \quad (2.13)$$

It is useful to keep several things in mind when coding this algorithm.

- The function  $F(l, r, E)$ , consisting of the energy, potential and centrifugal barrier, given in Eq. (2.10), is coded into a function  $F(L, R, E)$ , with  $L$  an integer and  $R$  and  $E$  being real variables.
- As you can see from Eq. (2.9a), the value of the wave function is needed for two values of the radial coordinate  $r$ , both beyond  $r_{\max}$ . We can take  $r_1$  equal to the first integration point beyond  $r_{\max}$  (if the grid constant  $h$  for the integration fits an integer number of times into  $r_{\max}$ , it is natural to take  $r_1 = r_{\max}$ ).  $r_2$  is larger than  $r_1$  and it is advisable to take it roughly half a wavelength beyond the latter. The wavelength is given by  $\lambda = 2\pi/k = 2\pi\hbar/\sqrt{2mE}$ . As both  $r_1$  and  $r_2$  are equal to an integer number times the integration step  $h$  – they will in general not differ exactly half a wavelength – the precise values of  $r_1$  and  $r_2$  are determined in the routine and output to the appropriate routine parameters.
- The starting value at  $r = 0$  is given by  $u(r = 0) = 0$ . We do not know the value of the derivative, which determines the normalisation of the resulting function – this normalisation can be determined afterward. We take  $u_l(0) = 0$  and  $u_l(h) = h^{l+1}$  ( $h$  is the integration step), which is the asymptotic approximation for  $u_l$  near the origin for a *regular* potential (for the H-Kr interaction potential which diverges strongly near the origin, we must use a different boundary condition as we shall see below).

– Programming exercise – Write a code for the Numerov algorithm. The input parameters to the routine must include the integration step  $h$ , the

radial quantum number  $l$ , the energy  $E$  and the radial coordinate  $r_{\max}$ ; on output it yields the coordinates  $r_1$  and  $r_2$  and the values of the wave function  $u_l(r_1)$  and  $u_l(r_2)$ .

When building a program of some complexity, it is very important to build it up step by step and to check every routine extensively. Comparison with analytical solutions is then of prime importance. We now describe several checks that should be performed after completion of the Numerov routine (it is also sensible to test a library routine).

**Check 1** The numerical solutions can be compared with analytical solutions for the case of the three-dimensional harmonic oscillator. Bound states occur for energies  $E = \hbar\omega(n + 3/2)$ ,  $n = 0, 1, 2, \dots$ . It is convenient in this case to choose units such that  $\hbar^2/2m = 1$ . Taking  $V(r) = r^2$ , we have  $\hbar\omega = 2$  and the lowest state occurs for  $l = 0$  with energy  $E = 3.0$ , with eigenfunction  $A \exp(-r^2/2)$ ,  $A$  being some constant. Using  $E = 3.0$  in our numerical integration routine should give us this solution with  $A = \exp(h^2/2)$  for the starting conditions described above. Check this for  $r$ -values up to  $r_2$ .

**Check 2** The integration method has an error of  $O(h^6)$  – the error found at the end of a finite interval then turns out to be less than  $O(h^4)$  (see problem A.3). This can be checked by comparing the numerical solution for the harmonic oscillator with the exact one. Carry out this comparison for several values of  $N$ , for example  $N = 4, 8, 16, \dots$ . For  $N$  large enough, the difference between the exact and the numerical solution should decrease for each new value of  $N$  by a factor of at least 16. If your program does not yield this behaviour, there must be an error in the code!

We shall now turn to the H-Kr interaction. The two-atom interaction potential for atoms is often modelled by the so-called Lennard-Jones (LJ) potential, which has the following form:

$$V_{\text{LJ}}(r) = \varepsilon \left[ \left( \frac{\sigma}{r} \right)^{12} - 2 \left( \frac{\sigma}{r} \right)^6 \right]. \quad (2.14)$$

This form of potential contains two parameters,  $\varepsilon$  and  $\sigma$ , and for H-Kr the best values for these are

$$\varepsilon = 5.9 \text{ meV} \quad \text{and} \quad \sigma = 3.57 \text{ \AA}. \quad (2.15)$$

Note that the energies are given in *milli-eV*! In units of meV and  $\sigma$  for energy and distance respectively, the factor  $2m/\hbar^2$  is equal to about  $6.12 \text{ meV}^{-1} \sigma^{-2}$ . The potential used by Toennies *et al.* included small corrections to the Lennard-Jones shape.

For the LJ potential the integration of the radial Schrödinger equation gives problems for small  $r$  because of the  $1/r^{12}$  divergence at the origin. We avoid integrating in this region and start at a nonzero radius  $r_{\min}$  where we use the analytic approximation of the solution for small  $r$  to find the starting values of the numerical solution. For  $r < r_{\min}$ , the term  $1/r^{12}$  dominates the other terms in the potential and the energy, so that the Schrödinger equation reduces to

$$\frac{d^2 u}{dr^2} = \epsilon \alpha \frac{1}{r^{12}} u(r) \quad (2.16)$$

with  $\alpha = 6.12$ . The solution of this equation is given by

$$u(r) = \exp(-Cr^{-5}) \quad (2.17)$$

with  $C = \sqrt{\epsilon \alpha / 25}$ . This fixes the starting values of the numerical solution at  $r_{\min}$  which should be chosen such that it can safely be assumed that the  $1/r^{12}$  dominates the remaining terms in the potential; typical values for the starting value of  $r$  lie between  $0.5\sigma$  and  $0.8\sigma$  (the minimum of the LJ potential is found at  $r = 2$ ). In problem 2.3 the starting solution for the LJ potential is considered in more detail.

You can adapt your program to the problem at hand by simply changing the function  $F(l, r, E)$  to contain the Lennard-Jones potential and by implementing the boundary conditions as described. As a check, you can verify that the solution does not become enormously large or remain very small.

### 2.2.2 The spherical Bessel functions

For the present problem, you need only the first six spherical Bessel functions  $j_l$  and  $n_l$ , and you can type in the explicit expressions directly. If you want a general routine for the spherical Bessel functions however, you can use the recursive procedures described in appendix A – see also problem A.1. Although upward recursion can be unstable for  $j_l$  (see appendix A), this is not noticeable for the small  $l$  values (up to  $l = 6$ ) which we need and you can safely use the simple upward recursion for both  $n_l$  and  $j_l$  (or use a library routine).

– Programming exercise – Write routines for generating the values of the spherical Bessel functions  $j_l$  and  $n_l$ . On input, the values of  $l$  and the argument  $x$  are specified and on output the value of the appropriate Bessel function is obtained.

and  $\sigma$  for energy  
at  $6.12 \text{ meV}^{-1} \sigma^{-2}$ .  
corrections to the

ger equation gives  
origin. We avoid  
where we use the  
starting values of  
the other terms  
tion reduces to

(2.16)

(2.17)

merical solution at  
med that the  $1/r^{12}$   
ues for the starting  
J potential is found  
ential is considered

mply changing the  
d by implementing  
can verify that the  
small.

al Bessel functions  
rectly. If you want  
ever, you can use  
also problem A.1.  
endix A), this is not  
d and you can safely  
a library routine).

ing the values of the  
values of  $l$  and the  
e appropriate Bessel

*Check* If your program is correct, it should yield the values for  $j_5$  and  $n_5$  given in problem A.1.

### 2.2.3 Putting the pieces together – results

To obtain the scattering cross sections, some extra routines must be added to the program. First of all, the phase shift must be extracted from the values  $r_1, u(r_1)$  and  $r_2, u(r_2)$ . This is straightforward using Eq. (2.9a). The total cross section can then readily be calculated using Eq. (2.8). The choice of  $r_{\max}$  must be made carefully, preferably keeping the error of the same order as the  $O(h^6)$  error of the Numerov routine (or the error of your library routine). In problem 2.2 it is shown that the deviation in the phase shift caused by cutting off the potential at  $r_{\max}$  is given by

$$\Delta\delta_l = -\frac{2m}{\hbar^2} k \int_{r_{\max}}^{\infty} j_l^2(kr) V_{\text{LJ}}(r) r^2 dr \quad (2.18)$$

and this formula can be used to estimate the resulting error in the phase shift or to improve the value found for it with a potential cut-off beyond  $r_{\max}$ . A good value is  $r_{\max} \approx 5\sigma$ .

For the determination of the differential cross section you will need additional routines for the Legendre polynomials.<sup>†</sup> In the following we shall only describe results for the total cross section.

– Programming exercise – *Add the necessary routines to the ones you have written so far and combine them into a program for calculating the total cross section.*

A computer program similar to the one described here was used by Toennies *et al.*<sup>7</sup> to compare the results of scattering experiments with theory. The experiment consisted of the bombardment of krypton atoms with hydrogen atoms. Figure 2.3 shows the Lennard–Jones interaction potential plus the centrifugal barrier  $l(l+1)/r^2$  of the radial Schrödinger equation. For higher  $l$ -values, the potential consists essentially of a hard core, a well and a barrier which is caused by the  $1/r^2$  centrifugal term in the Schrödinger equation. In such a potential, quasi-bound states are possible. These are states which would be genuine bound states for a potential for which the barrier does not drop to zero for larger values of  $r$ , but remains at its maximum height. You can imagine

<sup>†</sup>These can be generated using the recursion relation  $(l+1)P_{l+1}(x) = (2l+1)xP_l(x) - lP_{l-1}(x)$ .




Figure 2.3: The effective potential for the Lennard-Jones interaction for various  $l$ -values.

the following to happen when a particle is injected into the potential at precisely this energy: it tunnels through the barrier, remains in the well for a relatively long time, and then tunnels outward through the barrier in an arbitrary direction because it has 'forgotten' its original direction. In wave-like terms, the particle resonates in the well, and this state decays after a relatively long time. This phenomenon is called 'scattering resonance'. This means that particles injected at this energy are strongly scattered and this shows up as a peak in the total cross section.

Such peaks can be seen figure 2.4, which shows the total cross section as a function of the energy calculated with a program as described above. The peaks are due to  $l = 4$ ,  $l = 5$  and  $l = 6$  scattering, with energies increasing with  $l$ . Figure 2.5 finally shows the experimental results for the total cross section for H-Kr. We see that the agreement is excellent.

You should be able now to reproduce the data of figure 2.4 with your program.

### \*2.3 Calculation of scattering cross sections

In this section we derive Eqs. (2.7) and (2.8). At a large distance from the scattering centre we can make an *Ansatz* for the wave function. This consists

ic potential



eraction for various  $l$ -

potential at precisely  
well for a relatively  
an arbitrary direction  
ke terms, the particle  
vely long time. This  
that particles injected  
peak in the total cross

total cross section as  
described above. The  
ergies increasing with  
the total cross section

figure 2.4 with your

# sections

rge distance from the  
unction. This consists

## 2.3. Calculation of scattering cross sections




Figure 2.4: The total cross section shown as function of the energy for a Lennard-Jones potential modeling the H-Kr system. Peaks correspond to the resonant scattering states.

of the incoming beam and a scattered wave:

$$\psi(\mathbf{r}) \sim e^{i\mathbf{k}\cdot\mathbf{r}} + f(\theta) \frac{e^{ikr}}{r}. \quad (2.19)$$

$\theta$  is the angle between the incoming beam and the line passing through  $\mathbf{r}$  and the scattering centre.  $f$  does not depend on the azimuthal angle  $\phi$  because the incoming wave has azimuthal symmetry, and the spherically symmetric potential will not generate  $m \neq 0$  contributions to the scattered wave.  $f(\theta)$  is called the scattering amplitude. From the *Ansatz* it follows that the differential cross section is given directly by the square of this amplitude:

$$\frac{d\sigma}{d\Omega} = |f(\theta)|^2 \quad (2.20)$$

with the appropriate normalisation (see for example ref. 6).

Beyond  $r_{\max}$ , the solution can also be written in the form (2.2) leaving out all  $m \neq 0$  contributions because of the azimuthal symmetry:

$$\psi(\mathbf{r}) = \sum_{l=0}^{\infty} A_l \frac{u_l(r)}{r} P_l(\cos \theta) \quad (2.21)$$




Figure 2.5: Experimental results as obtained by Toennies *et al.*<sup>7</sup> for the total cross section (arbitrary units) of the scattering of hydrogen atoms by krypton atoms as function of centre of mass energy.

where we have used the fact that  $Y_0^l(\theta, \phi)$  is proportional to  $P_l[\cos(\theta)]$ . Because the potential vanishes in the region  $r > r_{\max}$ , the solution  $u_l(r)/r$  is given by the linear combination of the regular and irregular spherical Bessel functions, and as we have seen this reduces for large  $r$  to

$$u_l(r) \approx \sin(kr - \frac{l\pi}{2} + \delta_l). \quad (2.22)$$

We want to derive the scattering amplitude  $f(\theta)$  by equating the expressions (2.19) and (2.21) for the wave function. For large  $r$  we obtain, using (2.22):

$$\sum_{l=0}^{\infty} A_l \left[ \frac{\sin(kr - l\pi/2 + \delta_l)}{kr} \right] P_l(\cos \theta) = e^{i\mathbf{k} \cdot \mathbf{r}} + f(\theta) \frac{e^{ikr}}{r}. \quad (2.23)$$

We write the right hand side of this equation as an expansion similar to that in the left hand side, using the following expression for a plane wave<sup>8</sup>

$$e^{i\mathbf{k} \cdot \mathbf{r}} = \sum_{l=0}^{\infty} (2l+1) i^l j_l(kr) P_l(\cos \theta). \quad (2.24)$$

$f(\theta)$  can also be written as an expansion in Legendre polynomials:

$$f(\theta) = \sum_{l=0}^{\infty} f_l P_l(\cos \theta), \quad (2.25)$$

so that we obtain:

$$\sum_{l=0}^{\infty} A_l \left[ \frac{\sin(kr - l\pi/2 + \delta_l)}{kr} \right] P_l(\cos\theta) = \sum_{l=0}^{\infty} \left[ (2l+1) i^l j_l(kr) + f_l \frac{e^{ikr}}{r} \right] P_l(\cos\theta). \quad (2.26)$$

If we substitute the asymptotic form (2.5) of  $j_l$  in the right hand side, we find:

$$\sum_{l=0}^{\infty} A_l \left[ \frac{\sin(kr - l\pi/2 + \delta_l)}{kr} \right] P_l(\cos\theta) = \sum_{l=0}^{\infty} \left[ \frac{2l+1}{2ik} (-)^{l+1} e^{-ikr} + \left( f_l + \frac{2l+1}{2ik} \right) e^{ikr} \right] P_l(\cos\theta). \quad (2.27)$$

Both the left and the right hand side of (2.27) contain in- and outgoing spherical waves (the occurrence of incoming spherical waves does not violate causality: they arise from the incoming plane wave). For each  $l$ , the prefactors of the incoming and outgoing waves should both be equal on both sides in (2.27). This condition leads to

$$A_l = (2l+1) e^{i\delta_l} i^l \quad (2.28)$$

and

$$f_l = \frac{2l+1}{k} e^{i\delta_l} \sin(\delta_l). \quad (2.29)$$

Using (2.20), (2.25), and (2.29), we can write down an expression for the differential cross section in terms of the phase shifts  $\delta_l$ :

$$\frac{d\sigma}{d\Omega} = \frac{1}{k^2} \left| \sum_{l=0}^{\infty} (2l+1) e^{i\delta_l} \sin(\delta_l) P_l(\cos\theta) \right|^2. \quad (2.30)$$

For the total cross section we find, using the orthonormality relations of the Legendre polynomials:

$$\sigma_{\text{tot}} = 2\pi \int d\theta \sin\theta \frac{d\sigma}{d\Omega}(\theta) = \frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) \sin^2 \delta_l. \quad (2.31)$$